

II. SYNOPTIC CHARTS OF SOLAR MAGNETIC FIELDS

Mount Wilson Observatory

These Synoptic Charts are constructed from the digital data of the daily magnetograms obtained at the 150-foot Tower Telescope at Mount Wilson. The spectrum line employed is λ 5250.2, Fe I. The magnetograph at the Tower Telescope measures only the longitudinal component of the Zeeman effect. The aperture is 17 arc seconds on a side, and the whole area of the sun is generally covered in one magnetogram with no overlapping.

Each Synoptic Chart is made up of computerdrawn segments from individual day's observations. In order to get standardized charts covering one half of a solar rotation, the charts have been redrawn. There is no averaging done at any point from more than one day's observation.

The solid horizontal lines represent the equator, $\pm 20^\circ$, $\pm 40^\circ$, and $\pm 60^\circ$ latitude. This is an equal-area projection, so that the line at the top of the chart represents the north pole, and the line at the bottom of the chart represents the south pole.

The Carrington longitudes are given at the bottom of the chart. The solid vertical lines represent each even 10 degrees in longitude.

Straight horizontal dashed lines represent the dividing line between regions where there is data and regions where there is no data due to incomplete observations.

Vertical dashed lines represent the dividing lines between observations from different days.

The longitude of central meridian at the time of observation is indicated by a short vertical line and the data at the top of the chart.

Solid magnetic contour lines represent positive fields (magnetic vector pointed toward the observer), and dashed contour lines represent negative fields. Exceptions to this rule are made when a contour line represents a lower rather than a higher value. Thus if there is a positive 10 gauss contour line that is solid surrounding a dashed contour line, the latter represents positive 10 gauss, and within it is less than positive 10 gauss. This rule applies only to contours which are not open at a boundary.


The gauss levels are ± 5 , ± 10 , ± 20 , ± 40 , ± 80 gauss. Corrections are made to compensate for limb darkening or any other decrease in the brightness of the signal, but no corrections are made for possible geometrical effects such as the inclination of the line of sight to the lines of force at high latitudes.


Positions are generally accurate to about 15 arc seconds. Naturally there is some smearing of the data at high latitudes because of the geometry of the situation. Data greater than 40° from central meridian should be treated with some caution. Serious inaccuracies of fit at the boundary between two day's observations are normally due to rapid growth of a region.

ROBERT HOWARD

30.1.69

ROT. NO. 1543






27.2.69

17.2

ROT. NO. 1544



ROT. NO. 1544
1.2 311

10.269

9.2

8.2


4.2


3.2

2.2

1.2

311





12.3.69


7.3

6.3

5.3

4.3

ROT. NO. 1545



II. SYNOPTIC CHARTS OF SOLAR MAGNETIC FIELDS

Mount Wilson Observatory

These Synoptic Charts are constructed from the digital data of the daily magnetograms obtained at the 150-foot Tower Telescope at Mount Wilson. The spectrum line employed is λ 5250.2, Fe I. The magnetograph at the Tower Telescope measures only the longitudinal component of the Zeeman effect. The aperture is 17 arc seconds on a side, and the whole area of the sun is generally covered in one magnetogram with no overlapping.

Each Synoptic Chart is made up of computerdrawn segments from individual day's observations. In order to get standardized charts covering one half of a solar rotation, the charts have been redrawn. There is no averaging done at any point from more than one day's observation.

The solid horizontal lines represent the equator, $\pm 20^\circ$, $\pm 40^\circ$, and $\pm 60^\circ$ latitude. This is an equal-area projection, so that the line at the top of the chart represents the north pole, and the line at the bottom of the chart represents the south pole.

The Carrington longitudes are given at the bottom of the chart. The solid vertical lines represent each even 10 degrees in longitude.

Straight horizontal dashed lines represent the dividing line between regions where there is data and regions where there is no data due to incomplete observations.

Vertical dashed lines represent the dividing lines between observations from different days.

The longitude of central meridian at the time of observation is indicated by a short vertical line and the data at the top of the chart.

Solid magnetic contour lines represent positive fields (magnetic vector pointed toward the observer), and dashed contour lines represent negative fields. Exceptions to this rule are made when a contour line represents a lower rather than a higher value. Thus if there is a positive 10 gauss contour line that is solid surrounding a dashed contour line, the latter represents positive 10 gauss, and within it is less than positive 10 gauss. This rule applies only to contours which are not open at a boundary.

The gauss levels are ± 5 , ± 10 , ± 20 , ± 40 , ± 80 gauss. Corrections are made to compensate for limb darkening or any other decrease in the brightness of the signal, but no corrections are made for possible geometrical effects such as the inclination of the line of sight to the lines of force at high latitudes.

Positions are generally accurate to about 15 arc seconds. Naturally there is some smearing of the data at high latitudes because of the geometry of the situation. Data greater than 40° from central meridian should be treated with some caution. Serious inaccuracies of fit at the boundary between two day's observations are normally due to rapid growth of a region.

ROBERT HOWARD


8.4.69

1.4

31.3

28.3

ROT. NO. 1546



180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360

21.4.69

20.4


16.4

14.4

13.4


12.4

11.4



0 10 20 30 40 50 60 70 80 90 100 110 120 130 140 150 160 170 180

1.569 30.4 29.4 28.4



180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360

20.5.69

19.5

18.5

17.5

15.5


12.5


11.5

10.5

9.5

ROT. NO. 1547





16.6.69


14.5

12.6

7.6

5.6

ROT. NO. 1548



28.6.69

26.6


24.6

22.6


20.6

18.6

ROT. NO. 1549



180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360



II. SYNOPTIC CHARTS OF SOLAR MAGNETIC FIELDS

Mount Wilson Observatory

These Synoptic Charts are constructed from the digital data of the daily magnetograms obtained at the 150-foot Tower Telescope at Mount Wilson. The spectrum line employed is λ 5250.2, Fe I. The magnetograph at the Tower Telescope measures only the longitudinal component of the Zeeman effect. The aperture is a square, 17 arc seconds on a side, and the whole area of the sun is generally covered in one magnetogram with no overlapping.

Each Synoptic Chart is made up of computer-drawn segments from individual day's observations. In order to get standardized charts covering one half of a solar rotation, the charts have been redrawn. There is no averaging done at any point from more than one day's observation.

The solid horizontal lines represent the equator, $\pm 20^\circ$, $\pm 40^\circ$, and $\pm 60^\circ$ latitude. This is an equal-area projection, so that the line at the top of the chart represents the north pole, and the line at the bottom of the chart represents the south pole.

The Carrington longitudes are given at the bottom of the chart. The solid vertical lines represent each even 10 degrees in longitude.

Straight horizontal dashed lines represent the dividing line between regions where there is data and regions where there is no data due to incomplete observations.

Vertical dashed lines represent the dividing lines between observations from different days.

The longitude of central meridian at the time of observation is indicated by a short vertical line and the data at the top of the chart.


Solid magnetic contour lines represent positive fields (magnetic vector pointed toward the observer), and dashed contour lines represent negative fields. Exceptions to this rule are made when a contour line represents a lower rather than a higher value. Thus if there is a positive 10 gauss contour line that is solid surrounding a dashed contour line, the latter represents positive 10 gauss, and within it is less than positive 10 gauss. This rule applies only to contours which are not open at a boundary.


The gauss levels are ± 5 , ± 10 , ± 20 , ± 40 , ± 80 gauss. Corrections are made to compensate for limb darkening or any other decrease in the brightness of the signal, but no corrections are made for possible geometrical effects such as the inclination of the line of sight to the lines of force at high latitudes.


Positions are generally accurate to about 15 arc seconds. Naturally there is some smearing of the data at high latitudes because of the geometry of the situation. Data greater than 40° from central meridian should be treated with some caution. Serious inaccuracies of fit at the boundary between two day's observations are normally due to rapid growth of a region.


Partial support for the observation and reduction of this material comes from the U.S. Office of Naval Research through Contract No. N00014-66-C-0239.


ROBERT HOWARD











3.10.69

1.10

30.9

29.9

28.9

27.9

26.9

25.9


24.9

23.9

22.9

21.9

ROT. NO. 1552
20.9



II. SYNOPTIC CHARTS OF SOLAR MAGNETIC FIELDS

Mount Wilson Observatory

These Synoptic Charts are constructed from the digital data of the daily magnetograms obtained at the 150-foot Tower Telescope at Mount Wilson. The spectrum line employed is λ 5250.2, Fe I. The magnetograph at the Tower Telescope measures only the longitudinal component of the Zeeman effect. The aperture is a square, 17 arc seconds on a side, and the whole area of the sun is generally covered in one magnetogram with no overlapping.

Each Synoptic Chart is made up of computer-drawn segments from individual day's observations. In order to get standardized charts covering one half of a solar rotation, the charts have been redrawn. There is no averaging done at any point from more than one day's observation.

The solid horizontal lines represent the equator, $\pm 20^\circ$, $\pm 40^\circ$, and $\pm 60^\circ$ latitude. This is an equal-area projection, so that the line at the top of the chart represents the north pole, and the line at the bottom of the chart represents the south pole.

The Carrington longitudes are given at the bottom of the chart. The solid vertical lines represent each even 10 degrees in longitude.

Straight horizontal dashed lines represent the dividing line between regions where there is data and regions where there is no data due to incomplete observations.

Vertical dashed lines represent the dividing lines between observations from different days.

The longitude of central meridian at the time of observation is indicated by a short vertical line and the data at the top of the chart.

Solid magnetic contour lines represent positive fields (magnetic vector pointed toward the observer), and dashed contour lines represent negative fields. Exceptions to this rule are made when a contour line represents a lower rather than a higher value. Thus if there is a positive 10 gauss contour line that is solid surrounding a dashed contour line, the latter represents positive 10 gauss, and within it is less than positive 10 gauss. This rule applies only to contours which are not open at a boundary.


The gauss levels are ± 5 , ± 10 , ± 20 , ± 40 , ± 80 gauss. Corrections are made to compensate for limb darkening or any other decrease in the brightness of the signal, but no corrections are made for possible geometrical effects such as the inclination of the line of sight to the lines of force at high latitudes.


Positions are generally accurate to about 15 arc seconds. Naturally there is some smearing of the data at high latitudes because of the geometry of the situation. Data greater than 40° from central meridian should be treated with some caution. Serious inaccuracies of fit at the boundary between two day's observations are normally due to rapid growth of a region.

Partial support for the observation and reduction of this material comes from the U.S. Office of Naval Research through Contract No. N00014-66-C-0239.

ROBERT HOWARD

ROT. NO. 1553
4.10





13.11.69

12.11

5.11


4.11

3.11

2.11


1.11

ROT. NO. 1554
31.10



180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360

22.11.69 21.11 19.11 18.11 17.11



0 10 20 30 40 50 60 70 80 90 100 110 120 130 140 150 160 170 180


10.12.69

7.12

5.12


4.12

3.12



180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360

24.12.69 23.12 22.12 21.12 20.12 17.12 14.12 13.12 12.12



0 10 20 30 40 50 60 70 80 90 100 110 120 130 140 150 160 170 180